

#### LA-UR-21-27461

Approved for public release; distribution is unlimited.

Title: Planetary Defense Asteroids, Comets, Impact Craters, and Outwitting

the Dinosaurs

Author(s): Plesko, Catherine Suzanne

Intended for: Public talk to science summer camp

Issued: 2021-07-29



#### Planetary Defense

## Asteroids, Comets, Impact Craters, and Outwitting the Dinosaurs





Dr. Cathy Plesko

July 21, 2021




### Our busy solar system






Comets are more ice than rock.




Asteroids are more rock than ice.





## "All is number." -Pythagoras






Galen Gisler, Los Alamos National Laboratory

with T Heberling, C Plesko, R Weaver (LANL)

Results on water impacts of small (<300 m diameter) asteroids

- Impacts or airbursts near populated shores are dangerous.
- Impacts far from shore are spectacular and may produce detectable waves at great distances.
- Airbursts over water are less effective at generating waves than impacts into water.
- The most significant effect of an impact into an ocean is the injection of water vapor into the stratosphere, possibly affecting climate.


# Impact of a 250-meter diameter asteroid into a deep ocean




#### Volume rendering of density.

Visualizations in this presentation were made by a team at Los Alamos (John M. Patchett, David H. Rogers, Boonthanome Nouanesengsy) and University of Texas (Francesca G. Samsel, Karen Tsai, Gregory D. Abram, Terece L. Turton) using simulation data generated by Galen Gisler.

#### You don't want to be close to an impact site!




#### 250 m asteroid, no airburst, 45° trajectory





Volume rendering of water mass fraction (top) and pressure (bottom)


Direct comparison of 3 calculations showing pressure (green), asteroid mass fraction (red), water mass fraction (blue)




#### Calculations of different sized impactors







The wavelength of these surface waves is much smaller than typical tsunami wavelengths. They are therefore not expected to propagate efficiently over long distances.



#### Impact Prevention: Deflection

- Push to alter orbit
- Mechanisms:
  - okinetic
  - oexplosive
  - olight
  - ogravity
  - orockets
- Diameter > 100 m
- Reaction time ~ years to decades




#### **Preventing Impacts: Disrupt and Disperse**

- Fragment
- Scatter fragments
- Mechanisms:
  - kinetic
  - explosive
  - mechanical
- Works best for:
  - Shorter warning
  - Smaller objects



NASA: Deep Impact Mission




We can test the physics with an experiment

#### The Target: Asteroid Dimorphos

- Target of the DART kinetic impactor mission
- Smaller member of a binary system
- 100-m-diameter, rocky mixedtype asteroid, 25% porosity
- Tunguska Meteor-sized
- Modeled 4 kinetic impactor scenarios


Model	Impactor Mass [kg]	Impactor Velocity $[km/s]$	Impact Angle [°]	
1	10350.4	19.75	0	
2	10350.4	19.75	90	
3	7897.0	8.1	0	
4	7897.0	8.1	90	



#### **Computer Model Results**


- NASA designed trajectories assuming a perfectly inelastic collision (100% momentum transfer efficiency)
- We estimate 400-2000% momentum transfer efficiency
- Total system momentum is conserved
- Spall, shrapnel mean target mass is not constant
- So remaining target can have a higher momentum.


Model	$m_i$ [kg]	$v_i  [\mathrm{km/s}]$	$\theta_i$ [°]	$m_{ej}$ [kg]	β	$\Delta v \text{ [cm/s]}$
1	10350.4	19.75	0	$> 6.7 \times 10^7$	> 7	> 30
2	10350.4	19.75	90	$> 1.2 \times 10^9$	> 17	> 100
3	7897.0	8.1	0	$8.2 \times 10^{8}$	21	36
4	7897.0	8.1	90	$1.3 \times 10^8$	20	38



## Questions?

# How often do impacts happen, and how do we know that, exactly?





#### The Moon: No plate tectonics or liquid water.




#### Scientists count the craters



## The age of rocks collected on the Moon were estimated from radioactive decay of the minerals.



## Then we know how many there are of a given size, and how old they are.




If you know that, you can estimate how often objects of a given size hit the Moon (and the Earth).

But...


## Little asteroids hit less often than big ones, and all the big craters have been counted.



#### Fortunately, there's plenty of data



#### Some projects use crowdsourcing



#### Some projects use machine learning




Figure 2. MOC Image M0803054: Test Scene 1, re

## Modern automated counts still haven't beaten simple hand-programmed counting scripts



Clark Chapman wrote the first automated crater counting script more than 40 years ago.

